

Institute of Molecular Biology and Biotechnology (IMBB)

Foundation for Research and Technology- Hellas (FORTH)

PRESS RELEASE

Heraklion, Greece, June 28, 2019

IMBB researchers uncover a link between mitochondrial maturation and germ cell differentiation.

Unlike somatic cells, gametes are virtually immortal and fulfill the pivotal task of protecting and faithfully transferring genetic material across generations. A recent study published in the premier international scientific journal *Cell Death and Differentiation* (Nature Press), reveals that mitochondrial maturation and function is linked with germ cell differentiation in the nematode *Caenorhabditis elegans*. Mitochondria are cellular organelles of prokaryotic origin which execute numerous functions in eukaryotic cells, with ATP production being the most prominent one.

IMBB researchers Dr. Nikolaos Charmpilas and Dr. Nektarios Tavernarakis (Professor at the Medical School, University of Crete, and Chairman of the Board at FORTH) demonstrated that inhibition of mitochondrial genome (mtDNA) transcription leads to germline tumor formation in nematodes, due to impaired differentiation and oocyte production. Thorough monitoring of germline mitochondria highlighted that they transition from a globular to a tubular shape during differentiation. Elongated organelles, in contrast with globular mitochondria of undifferentiated germ cells, exhibit various signs of functional maturation, such as, increased ATP production and elevated membrane potential, as well as, enhanced production of reactive oxygen species (ROS). This mitochondrial maturation process is tightly controlled by sperm-originating (MSP) signals, which are known to promote oogenesis and ovulation.

These findings integrate mitochondria and their homeostasis in the developmental modules which shape the *C. elegans* germline. Moreover, establish enhanced mitochondrial maturation as a prerequisite for stem cell differentiation in diverse model organisms, and further implicate augmented mitochondrial activity in tumorigenesis.

For additional information, please contact:


Prof. Nektarios Tavernarakis
Chairman of the Board, FORTH
Research Director, IMBB-FORTH
Professor, Medical School, University of Crete
eMail: tavernarakis@imbb.forth.gr

Relevant links:

www.elegans.gr

<https://www.nature.com/cdd/>

<https://www.nature.com/articles/s41418-019-0375-9>



Globular mitochondria
 Low ATP production
 Low mitoROS
 Low membrane potential

Elongated mitochondria
 High ATP production
 High mitoROS
 High membrane potential

